Instructions for Use



Silvet® is a hypertonic buffered solution, with a pleasant odor, for wound disinfection, management and maintenance of wound moisture level. Silvet is a clear, colorless solution, protected from microbial contamination by a low concentration of ionic silver and supplied in 100mL, 250mL and 500mL bottles.

INTENDED USE AND INDICATIONS FOR USE

Silvet® is intended for use in animals and pets, by or under the supervision of a veterinarian for disinfection, management, moisturizing and healing acceleration of wounds, different types of ulcers, post-surgical wounds, first and second-degree burns, cuts abrasions and minor skin irritations.

GENERAL DIRECTIONS FOR USE

Silvet® is designed for management and moisturizing of wounds and burns. Facilitate wound healing by applying the **Silvet®** solution directly from the bottle by means of a syringe or bulb, spraying onto the wound using a spray bottle, by applying with devices designed to allow for irrigation or lavage, or by soaking a gauze/dressing or sponge.

Rinse the wound with **Silvet®** and clean using gauze. Soak dry gauze with **Silvet®** solution and dress the wound. The soaked gauze can be covered with semi-occlusive dressing in order to prevent solution evaporation.

Apply 1-2 times daily, or every second day, according to the veterinarian's instruction.

Silvet® solution is ideal for in house treatment of wounds and serves as an integral part of a first aid kit for treatment of pets and other animals. Use up to 10mL of **Silvet®** solution per each Kg body weight in 24 hours.

Silvet® should be used by itself as a stand-alone solution. The wound can be washed with water or soapy water prior to the application. The use of any other solutions in combination with SilVet® might reduce its efficacy.

Silvet® solution can be applied on the wound via a syringe, a spray bottle, irrigation, lavage system or pouring onto a gauze/dressing or sponge. **DO NOT INJECT**.

Note: The wound may be cleansed if Silvet® is applied with a 19-22G needle (application pressure of ~10 psi).



CAUTIONS AND WARNINGS

- For external use only. Avoid contact with eyes.
- Use only undamaged bottle.
- Discontinue use if irritation develops.
- Avoid usage of turbid or colored solution.
- Protect bottle from direct contact with the wound.
- Bottle that comes in direct contact with the wound must be discarded after use.
- For single-patient use only.
- Store between 10˚C-30°C (50˚F-86°F) in a dry place.
- If the bottle is closed immediately after use and the top of the bottle is protected from contamination, the solution may be reused within 28 days from first opening.
- The empty bottle can be disposed to the regular trash can. A bottle that contains solution should be cleared off to an appropriate site.

DIRECTIONS FOR USE

Step 1:



Open the cap, insert syringe with needle into the bottle and draw up the quantity of Silvet® solution as advised by the Medical Practitioner.

Step 2:



Rinse, by spraying the Silvet® solution on the wound. Let the wound be exposed to the sprayed solution for 5-10 minutes **DO NOT INJECT** (The needle should not touch the wound)

Step 3:



Wet a sterile cotton gauze with Silvet® solution.

Step 4:



Place the wet gauze on the wound and apply a secondary non-antimicrobial occlusive dressing as directed by the Medical Practitioner.

Step 5:



Before removal of the gauze/dressing or sponge, moisten with Silvet® solution for easier removal.

SilVet® solution can be applied on the wound via a syringe, a spray bottle, irrigation, lavage system, or pouring onto a gauze/dressing or sponge.

Notes:

- DO NOT wash the wound with Saline, Before or After using SilVet®.
- Do not use together with chloride ions containing fluids. Chloride ions bind to and inactivate silver ions, significantly reducing Silvet efficacy.

INGREDIENTS

Ingredients include water for injection (WFI), glycerol (glycerin), sorbitan monolaurate (Tween-20), TRIS, menthol and silver nitrate 0.01% (w/v).

Each ml contains ionic silver in the form of silver nitrate 0.01% (w/v).

Each package contains nine (12) bottles of 500 mL or 250mL, or twenty four (24) bottles of 100mL and instructions for use leaflet, which is part of the bottle label.

SYMBOLS

The following provides a description of the graphic symbols that appear on the SilVet label:

	Caution
	Consult instructions for use
	Keep bottles out of direct sunlight
	Temperature Limitation
	Single patient -multiple use
	Sterilized using aseptic processing techniques
	Do not resterilize. Resterilization may reduce the effectiveness of the product
	Do not use if package is damaged
	Keep out of reach of children
	19G needle (application pressure of ~ 10 psi)
MD	Medical device
LOT	Batch code / LOT #
	Date of manufacture Country of manufacture
	Use by date
REF	Catalog number
	Manufacturer
	Distributor
UDI	UDI - Unique device identifier



SilverStream Medical Ltd.
Company Reg. No. 513160986
26 Shabazi St., Rosh Ha'Ayin 4802126, Israel
Phone: +972 3 9194334
WWW.SIL.VET